

Quantum mechanics on rational numbers

A. Vourdas
University of Bradford

- background
 - $\mathbb{Q}_p/\mathbb{Z}_p$ and its Pontryagin dual group \mathbb{Z}_p
 - \mathbb{Q}/\mathbb{Z} and its Pontryagin dual group $\widehat{\mathbb{Z}}$
 - \mathbb{Q} and its Pontryagin dual group $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$
 - $\mathbb{Q}^{(\pi)}$ and its Pontryagin dual \mathbb{S}_{π} (solenoid)
 - $n^{-1}\mathbb{Z}$ and its Pontryagin dual $\mathbb{R}/n\mathbb{Z}$
- The Schwartz-Bruhat space for $S[(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}), \mathbb{Q}]$
- examples: $S(\mathbb{S}_{\pi}, \mathbb{Q}^{(\pi)})$, $S[(\mathbb{R}/n\mathbb{Z}), n^{-1}\mathbb{Z}]$
- Heisenberg-Weyl group and other phase space methods in $(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}) \times \mathbb{Q}$
- the set of subsystems of $S[(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}), \mathbb{Q}]$
 - partial order
 - T_0 -topology
- Discussion

A. Vourdas, JMAA 394, 48 (2012)

$\mathbb{Q}_p/\mathbb{Z}_p$ and its Pontryagin dual group \mathbb{Z}_p

- \mathbb{Z}_p as inverse limit.

$$\varprojlim \mathbb{Z}(p^\ell) = \mathbb{Z}_p$$

\mathbb{Z}_p profinite: compact, totally disconnected group

$$a_p = \bar{a}_0 + \bar{a}_1 p + \dots; \quad 0 \leq \bar{a}_i \leq p - 1$$

- Pontryagin dual group of \mathbb{Z}_p is $\mathbb{Q}_p/\mathbb{Z}_p$
 $\mathbb{Q}_p/\mathbb{Z}_p$ as direct limit.

$$\varinjlim \mathbb{Z}(p^\ell) = \mathbb{Q}_p/\mathbb{Z}_p$$

fractional p -adic numbers (cosets)

$$\mathfrak{b}_p = \bar{\mathfrak{b}}_{-k} p^{-k} + \dots + \bar{\mathfrak{b}}_{-1} p^{-1}$$

$\mathbb{Q}_p/\mathbb{Z}_p$ isomorphic to Prüfer group $\mathcal{C}(p^\infty)$

- characters in $\mathbb{Q}_p/\mathbb{Z}_p$

$$\chi_p(a_p \mathfrak{b}_p) = \exp(i2\pi a_p \mathfrak{b}_p)$$

\mathbb{Q}/\mathbb{Z} and its Pontryagin dual group $\widehat{\mathbb{Z}}$

- $\widehat{\mathbb{Z}}$ as inverse limit

$$\lim_{\leftarrow} \mathbb{Z}(\ell) = \widehat{\mathbb{Z}}$$

$\widehat{\mathbb{Z}}$ profinite group

$$\widehat{\mathbb{Z}} = \prod_{p \in \Pi} \mathbb{Z}_p$$

elements

$$s = (s_2, \dots, s_p, \dots); \quad s_p \in \mathbb{Z}_p; \quad p \in \Pi$$

\mathbb{Z} is embedded into $\widehat{\mathbb{Z}}$:

$$\mathbb{Z} \ni n \rightarrow (n, n, n, \dots) \in \widehat{\mathbb{Z}}$$

- Pontryagin dual group of $\widehat{\mathbb{Z}}$ is \mathbb{Q}/\mathbb{Z}
 \mathbb{Q}/\mathbb{Z} as direct limit:

$$\lim_{\rightarrow} \mathbb{Z}(\ell) = \mathbb{Q}/\mathbb{Z} \quad \mathbb{Q}/\mathbb{Z} = \bigoplus_{p \in \Pi} \mathbb{Q}_p/\mathbb{Z}_p$$

$x \in \mathbb{Q}/\mathbb{Z}$: (x_2, \dots, x_p, \dots) , where $x_p \in \mathbb{Q}_p/\mathbb{Z}_p$, $p \in \Pi$
all but a finite number of the x_p zero

- characters in \mathbb{Q}/\mathbb{Z}

$$\chi(sx) = \prod_{p \in \Pi} \chi_p(s_p x_p)$$

converges: only a finite number of the $\chi_p(s_p x_p)$ different from 1.

\mathbb{Q} and its Pontryagin dual group $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$

- ring $\mathbb{A}_{\mathbb{Q}}$ of adeles

$$y = (y_{\infty}, y_2, \dots, y_p, \dots); \quad y_p \in \mathbb{Q}_p; \quad \mathbb{Q}_{\infty} = \mathbb{R}$$

but $y_p \in \mathbb{Z}_p$ for all but a finite number of p
 $\mathbb{A}_{\mathbb{Q}}$: **restricted** direct product of \mathbb{Q}_p with respect to \mathbb{Z}_p :

$$\mathbb{A}_{\mathbb{Q}} = \prod' \mathbb{Q}_p$$

- \mathbb{Q} embedded into $\mathbb{A}_{\mathbb{Q}}$ as

$$\mathbb{Q} \ni u \rightarrow (u, u, u, \dots) \in \mathbb{A}_{\mathbb{Q}}$$

- $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$ as inverse limit

$$\lim_{\leftarrow} \mathbb{R}/n\mathbb{Z} \cong \mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$$

$\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$ describes finite covers of S .

- Pontryagin dual group of $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$ is \mathbb{Q}
 \mathbb{Q} as direct limit

$$\lim_{\rightarrow} n^{-1}\mathbb{Z} \cong \mathbb{Q}$$

- Additive characters on \mathbb{Q} are given by

$$\psi(uy) = \exp[i2\pi(-uy_\infty + uy_2 + \dots)] = \prod_{p \in \Pi_\infty} \chi_p(uy_p)$$

$$u \in \mathbb{Q}; \quad y \in \mathbb{A}_\mathbb{Q}/\mathbb{Q}$$

$$\chi_\infty(uy_\infty) = \exp(-i2\pi uy_\infty)$$

minus sign in $\chi_\infty(uy_\infty)$ and a plus sign in $\chi_p(uy_p)$

$y \rightarrow y + \text{rational}$: same result

convergence: finite number of factors $\neq 1$

- fundamental domain for $\mathbb{A}_\mathbb{Q}/\mathbb{Q}$ is $\mathbb{S} \times \widehat{\mathbb{Z}}$
($\mathbb{S} = \mathbb{R}/\mathbb{Z}$).

- $\mathbb{Q}^{(\pi)}$: subgroup of \mathbb{Q}

$$\mathbb{Q}^{(\pi)} = \left\{ \frac{a}{\prod_{p_i \in \pi} p_i^{e_i}} \mid a \in \mathbb{Z} \right\}$$

Pontryagin dual group of $\mathbb{Q}^{(\pi)}$: solenoid \mathbb{S}_π
 $(y_0, y_2, \dots, y_p, \dots)$ where $y_p = 0$ for $p \notin \pi$

$$\varprojlim \mathbb{R}/k^n \mathbb{Z} \cong \mathbb{S}_\pi; \quad k = \prod_{p \in \pi} p^e$$

special case: $\pi = \{p\}$:
 p -adic solenoid \mathbb{S}_p with elements (y_0, y_p) .

- subgroup of $\mathbb{Q}^{(\pi)}$

$$n^{-1}\mathbb{Z} = \left\{ \frac{a}{n} = \frac{a}{\prod_{p_i \in \pi} p_i^{e_i}} \mid e_i \leq E_i \right\}; \quad n = \prod_{p \in \pi} p_i^{E_i}$$

Pontryagin dual group $\mathbb{R}/n\mathbb{Z}$

elements of $\mathbb{R}/n\mathbb{Z}$: (y_0, k) where $y_0 \in \mathbb{S}$ and
 $k \in \mathbb{Z}(n)$ winding number

$$\mathbb{Z}(n) \cong \prod_{p \in \pi} \mathbb{Z}(p_i^{E_i})$$

chinese remainder theorem: winding number
 $k = (y_{p_1}, \dots)$ where $y_{p_i} \in \mathbb{Z}(p_i^{E_i})$

elements of $\mathbb{R}/n\mathbb{Z}$: $(y_0, y_2, \dots, y_p, \dots)$

$y_{p_i} \in \mathbb{Z}(p_i^{E_i})$ component of winding number for
that prime

The Schwartz-Bruhat space for $S[(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}), \mathbb{Q}]$

first: space Σ functions on $\mathbb{A}_{\mathbb{Q}}$

later: space \mathfrak{S} functions on $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$

- Schwartz-Bruhat space Σ : finite linear combinations of complex functions $\phi(y)$

$$\phi(y) = \phi_{\infty}(y_{\infty}) \prod_{p \in \Pi} \phi_p(y_p); \quad y = (y_{\infty}, y_2, \dots, y_p, \dots) \in \mathbb{A}_{\mathbb{Q}}$$

where

- (1) $\phi_{\infty}(y_{\infty}) \in \mathcal{S}(\mathbb{R})$ (Schwartz)
- (2) $\phi_p(y_p)$ locally constant complex functions with compact support,
- (3) for all but a finite number of $p \in \Pi$
 $\phi_p(y_p) = 1$ if y_p is p-adic integer

$\Pi[\phi(y)]$ contains indices $\phi_p(y_p) \neq 1$

$\Pi_1[\phi(y)]$ subset $y_p \in \mathbb{Q}_p$ (finite)

$\Pi_2[\phi(y)]$ subset $y_p \in \mathbb{Z}_p$ (finite)

- Integrals of functions in Σ over $\mathbb{A}_{\mathbb{Q}}$:

$$\begin{aligned} \int_{\mathbb{A}_{\mathbb{Q}}} \phi(y) dy &= \int_{\mathbb{R}} \phi_{\infty}(y_{\infty}) dy_{\infty} \\ &\times \prod_{p \in \Pi_1[\phi(y)]} \int_{\mathbb{Q}_p} \phi_p(y_p) dy_p \prod_{p \in \Pi_2[\phi(y)]} \int_{\mathbb{Z}_p} \phi_p(y_p) dy_p \end{aligned}$$

finite number of factors $\neq 1$

p-adic integrals of locally constant functions with constant support = finite sums.

- space \mathfrak{S}

From $\phi(y)$ on $\mathbb{A}_{\mathbb{Q}}$, to $f(\eta)$ on $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$

Weil transf: add values of $\phi(y)$ in each coset in $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$:

$$f(\eta) = \int_{\mathbb{Q}} du \phi(y + u); \quad \eta = \{y + u \mid u \in \mathbb{Q}\}; \quad \eta \in \mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$$

then

$$\int_{\mathbb{A}_{\mathbb{Q}}} \phi(y) dy = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} f(\eta) d\eta$$

- Fourier transform of $f(\eta)$

$$F(u) = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} f(\eta) \psi(u\eta) d\eta = \int_{\mathbb{A}_{\mathbb{Q}}} \phi(y) \psi(yu) dy; \quad u \in \mathbb{Q}$$

write $F(u)$ as

$$F(u) = F_{\infty}(u) \prod_{p \in \Pi[\phi(y)]} F_p(u) \prod_{p \notin \Pi[\phi(y)]} \Delta_p(u)$$

Fourier trans. of $\phi_p(y_p) = 1, y_p \in \mathbb{Z}_p$:

$$\Delta_p(u) = 0 \text{ if } u \neq 0$$

$$\Delta_p(0) = 1 \text{ zero coset in } \mathbb{Q}_p/\mathbb{Z}_p: \text{ p-adic integers}$$

$u = a/b$ is p-adic integer if $p \nmid b$

$$\Delta_p(u) = 1 \text{ if } u = a/b \text{ with } p \nmid b$$

- inverse Fourier transform

$$f(\eta) = \int_{\mathbb{Q}} du F(u) \psi(-u\eta); \quad \eta \in \mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$$

- scalar product

$$(f, g) = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} \overline{f(\eta)} g(\eta) d\eta; \quad (F, G) = \int_{\mathbb{Q}} du \overline{F(u)} G(u)$$

- Parseval: $(f, g) = (F, G)$

below examples with variables in subgroups of \mathbb{Q}
and in their Pontryagin dual groups

work in fundamental domain $\mathbb{S} \times \widehat{\mathbb{Z}}$ of $\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}$.

Examples

- QM for $S[(\mathbb{R}/\mathbb{Z}), \mathbb{Z}]$
 $f(\eta) = f_\infty(\eta_\infty)$ where $\eta_\infty \in \mathbb{S}$
 (i.e., $f_p(\eta_p) = 1$ for all $p \in \Pi$)
 Fourier transform

$$\begin{aligned}
 F(u) &= \int_{\mathbb{S} \times \widehat{\mathbb{Z}}} f_\infty(\eta_\infty) \psi(u\eta) d\eta \\
 &= \left[\int_0^1 d\eta_\infty f_\infty(\eta_\infty) \chi_\infty(u\eta_\infty) \right] \\
 &\quad \times \prod_{p \in \Pi} \Delta_p(u)
 \end{aligned}$$

$F(u)$ non-zero: if $u \in \mathbb{Z}$
 i.e., $u = a/b$ with $p \nmid b$ for **all** primes

- QM for $S(\mathbb{S}_{p_1}, \mathbb{Q}^{(p_1)})$
 $f(\eta) = f_\infty(\eta_\infty) f_{p_1}(\eta_{p_1})$
 (i.e., $f_p(\eta_p) = 1$ for all $p \in \Pi - \{p_1\}$)
 Fourier transform

$$\begin{aligned}
 F(u) &= \int_{\mathbb{S} \times \widehat{\mathbb{Z}}} f_\infty(\eta_\infty) f_{p_1}(\eta_{p_1}) \psi(u\eta) d\eta \\
 &= \left[\int_0^1 d\eta_\infty f_\infty(\eta_\infty) \chi_\infty(u\eta_\infty) \int_{\mathbb{Z}_{p_1}} d\eta_{p_1} f_{p_1}(\eta_{p_1}) \chi_{p_1}(u\eta_{p_1}) \right] \\
 &\quad \times \prod_{p \in \Pi - \{p_1\}} \Delta_p(u)
 \end{aligned}$$

$F(u)$ non-zero only if $u \in \mathbb{Q}^{(p_1)}$
 i.e., $u = a/b$ with $p \nmid b$ for $p \in \Pi - \{p_1\}$

- QM for $S[(\mathbb{R}/p_1^{e_1}\mathbb{Z}), p_1^{-e_1}\mathbb{Z}]$
 restrict the above formalism further
 $f_{p_1}(\eta_{p_1})$ locally constant with **given degree** e_1 :
 $f_{p_1}(\eta_{p_1} + \mathfrak{a}_{p_1}) = f_{p_1}(\eta_{p_1})$ for $|\mathfrak{a}_{p_1}|_{p_1} \leq p_1^{-e_1}$

η : pair $(\eta_\infty, \eta_{p_1})$ with $\eta_\infty \in \mathbb{R}/\mathbb{Z}$ and $\eta_{p_1} \in \mathbb{Z}(p_1^{e_1})$
 describes points in circle $\mathbb{R}/(p_1^{e_1}\mathbb{Z})$ wrapped $p_1^{e_1}$
 times around the circle \mathbb{R}/\mathbb{Z}

- QM for $S(\mathbb{S}_\pi, \mathbb{Q}^{(\pi)})$ with $\pi = \{p_1, \dots, p_\ell\}$

$$f(\eta) = f_\infty(\eta_\infty) f_{p_1}(\eta_{p_1}) \dots f_{p_\ell}(\eta_{p_\ell})$$

Fourier transform $F(u)$ non-zero only if
 $u \in \mathbb{Q}^{(\pi)}$

- QM for $S[(\mathbb{R}/n\mathbb{Z}), n^{-1}\mathbb{Z}]$ with $n = p_1^{e_1} \dots p_\ell^{e_\ell}$
 restrict the formalism further
 $f_{p_j}(\eta_{p_j})$, locally constant, with given degrees e_j

Phase space methods in $(\mathbb{A}_\mathbb{Q}/\mathbb{Q}) \times \mathbb{Q}$

- The Heisenberg-Weyl group $\mathcal{HW}(\mathbb{A}_\mathbb{Q}/\mathbb{Q}, \mathbb{Q}, \mathbb{A}_\mathbb{Q}/\mathbb{Q})$ displacement operators $D(\mathbf{a}, b, \mathbf{c})$

$$\begin{aligned} [\mathcal{D}(\mathbf{a}, b, \mathbf{c})F](u) &= \psi(\mathbf{c} - \mathbf{a}b + 2\mathbf{a}u) F(u - b) \\ [\mathcal{D}(\mathbf{a}, b, \mathbf{c})f](\eta) &= \psi(\mathbf{c} + \mathbf{a}b - \eta b) f(\eta - 2\mathbf{a}) \\ \mathbf{a}, \mathbf{c}, \eta &\in \mathbb{A}_\mathbb{Q}/\mathbb{Q}; \quad b, u \in \mathbb{Q} \end{aligned}$$

$\mathcal{HW}(\mathbb{A}_\mathbb{Q}/\mathbb{Q}, \mathbb{Q}, \mathbb{A}_\mathbb{Q}/\mathbb{Q})$ locally compact topological group.

- For any trace class operator θ

$$\begin{aligned} \int_{\mathbb{A}_\mathbb{Q}/\mathbb{Q}} d\mathbf{a} \int_{\mathbb{Q}} db \mathcal{D}(\mathbf{a}, b, 0) \theta [\mathcal{D}(\mathbf{a}, b, 0)]^\dagger &= \mathbf{1} \text{tr} \theta \\ \theta &= \int_{\mathbb{A}_\mathbb{Q}/\mathbb{Q}} d\mathbf{a} \int_{\mathbb{Q}} db \mathcal{D}(\mathbf{a}, b, 0) \text{tr}[\theta \mathcal{D}(-\mathbf{a}, -b, 0)] \end{aligned}$$

- coherent states

$$f_{\text{coh}}(\eta|\mathbf{a}, b) \equiv [\mathcal{D}(\mathbf{a}, b, \mathbf{c})f](\eta); \quad \mathbf{a} \in \mathbb{A}_\mathbb{Q}/\mathbb{Q}; \quad b \in \mathbb{Q}$$

with resolution of the identity:

$$\int_{\mathbb{A}_\mathbb{Q}/\mathbb{Q}} d\mathbf{a} \int_{\mathbb{Q}} db f_{\text{coh}}(\eta|\mathbf{a}, b) \overline{f_{\text{coh}}(\eta'|\mathbf{a}, b)} = \delta_A(\eta - \eta')$$

- parity around origin: $\mathcal{P}(0, 0)F(u) = F(-u)$
 parity around $(\mathbf{a}, b) \in (\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}) \times \mathbb{Q}$

$$\mathcal{P}(\mathbf{a}, b) = \mathcal{D}(\mathbf{a}, b, c) \mathcal{P}(0, 0) [\mathcal{D}(\mathbf{a}, b, c)]^\dagger$$

Parity related to displacements with Fourier tr

$$\mathcal{P}(\mathbf{a}, b) = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} d\mathbf{a}' \int_{\mathbb{Q}} db' \mathcal{D}(\mathbf{a}', b', 0) \psi(2\mathbf{a}'b - 2ab')$$

operator θ can be expanded as

$$\theta = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} d\mathbf{a} \int_{\mathbb{Q}} db \mathcal{P}(\mathbf{a}, b,) \text{tr}[\theta \mathcal{P}(\mathbf{a}, b)]$$

- Given a pair of functions $g(\eta), f(\eta) \in \mathfrak{S}$
 Wigner $\mathcal{W}(\mathbf{a}, b; g, f) = (g, \mathcal{P}(\mathbf{a}, b)f)$
 Weyl (or ambiguity) $\widetilde{\mathcal{W}}(\mathbf{a}, b; g, f) = (g, \mathcal{D}(\mathbf{a}, b, 0)f)$

$$\mathcal{W}(\mathbf{a}, b; g, f) = \int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} d\mathbf{a}' \int_{\mathbb{Q}} db' \widetilde{\mathcal{W}}(\mathbf{a}', b'; g, f) \psi(2\mathbf{a}'b - 2ab')$$

'marginal properties' of Wigner function

$$\int_{\mathbb{Q}} db \mathcal{W}(\mathbf{a}, b; g, f) = \overline{g(-2\mathbf{a})} f(-2\mathbf{a})$$

$$\int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} d\mathbf{a} \mathcal{W}(\mathbf{a}, b; g, f) = \overline{\widetilde{g}(-b)} \widetilde{f}(-b)$$

'marginal properties' of Weyl function:

$$\int_{\mathbb{Q}} db \widetilde{\mathcal{W}}(\mathbf{a}, b; g, f) = \overline{g(\mathbf{a})} f(-\mathbf{a})$$

$$\int_{\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}} d\mathbf{a} \widetilde{\mathcal{W}}(\mathbf{a}, b; g, f) = \overline{\widetilde{g}(2^{-1}b)} \widetilde{f}(-2^{-1}b)$$

the set of subsystems of $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$

- supernatural (Steinitz) numbers:

$$\mathbb{N}_S = \left\{ \prod p^{e_p} \mid p \in \Pi; \quad e_p \in \mathbb{Z}_0^+ \cup \{\infty\} \right\}$$

examples:

$$\Omega = \prod_{p \in \Pi} p^\infty; \quad \Omega(\pi) = \prod_{p \in \pi} p^\infty; \quad \Omega(\pi) | \Omega$$

directed partially ordered set

with divisibility as order: $m \prec n$ means $m|n$

Ω 'top element'

- directed-complete partial order (dcpo): each chain has a supremum

\mathbb{N}_S : dcpo

$p, p^2, p^3, \dots, p^\infty$ has p^∞ as supremum

$m, m^2, \dots, \Omega[\pi(m)]$ has $\Omega[\pi(m)]$ as supremum

- Topological space $(\mathbb{N}_S, T_{\mathbb{N}_S})$ with the 'divisor topology' $T_{\mathbb{N}_S}$ generated by the base

$$B_{\mathbb{N}_S} = \{\emptyset, U(n) \mid n \in \mathbb{N}_S\}; \quad U(n) = \{m \in \mathbb{N}_S \mid m|n\}$$

T_0 topological space (but not T_1)

separation axioms: T_2 (Hausdorff) $\subset T_1 \subset T_0$

- partial order: \mathbb{N}_S : dcpo, \mathbb{N} not dcpo
topology: \mathbb{N}_S : compact, \mathbb{N} locally compact, not compact

\mathbb{N} : something is missing!!

\mathbb{N}_S : we found what was missing!!

- subsystem: $S(E, \tilde{E}) \prec S(G, \tilde{G})$ if \tilde{E} subgroup of \tilde{G}
then quotient relation between E, G with annihilators

- A_S : set of subsystems of $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$
bijective map between A_S and \mathbb{N}_S :
 $S[(\mathbb{R}/n\mathbb{Z}), n^{-1}\mathbb{Z}] \leftrightarrow n \in \mathbb{N}$
 $S(\mathbb{S}_\pi, \mathbb{Q}^{(\pi)}) \leftrightarrow \Omega(\pi)$
 $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}] \leftrightarrow \Omega$

also $A = \{S[(\mathbb{R}/n\mathbb{Z}), n^{-1}\mathbb{Z}] \mid n \in \mathbb{N}\}$
bijective map between A and \mathbb{N}

- A_S order isomorphic to \mathbb{N}_S

embeddings of subsystems into supersystems
and their **compatibility**:
quantum states, Heisenberg-Weyl groups, Wigner functions, etc

$S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$ maximum element
 $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$: QM on 'large circles'

A not dcpo (something is missing!!)
 A_S dcpo (we found what was missing!!)

- (A_S, T_{A_S}) topological space homeomorphic to $(\mathbb{N}_S, T_{\mathbb{N}_S})$

T_0 -topology:

axioms of T_0 topology express basic logical relations between subsystems and supersystems

can be used to define continuity of a quantity (eg entropy) in systems and their supersystems

- A locally compact, not compact (something is missing!!)
 A_S compact (we found what was missing!!)
- $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$ smallest system that contains all $S(\mathbb{S}_\pi, \mathbb{Q}^{(\pi)})$, $S[(\mathbb{R}/q\mathbb{Z}), q^{-1}\mathbb{Z}]$ as subsystems

details for $S[(\mathbb{A}_\mathbb{Q}/\mathbb{Q}), \mathbb{Q}]$: work in progress

details for $S[\widehat{\mathbb{Z}}, (\mathbb{Q}/\mathbb{Z})]$:

A. Vourdas JMP53, 122101 (2012)

A. Vourdas JPA46 (2013) 043001

Discussion

- The Schwartz-Bruhat space for $S[(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}), \mathbb{Q}]$
- examples: $S(S_{\pi}, \mathbb{Q}^{(\pi)})$, $S[(\mathbb{R}/n\mathbb{Z}), n^{-1}\mathbb{Z}]$
- Heisenberg-Weyl group and other phase space methods in $(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}) \times \mathbb{Q}$
- partial order and topology of the set of sub-systems of $S[(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}), \mathbb{Q}]$
directed complete partial order
topology: compact
QM on large circles $\mathbb{R}/n\mathbb{Z}$